PRODUCT GUIDE / COLLECTIONS



Your Journey To Regenerative Health NATURALLY Starts With RegenX Naturals.



LSF-2 by RegenX Naturals

Immune System Support

Product Details

Lyme Support Formula 2 is used in cases where the primary Lyme symptoms may also have some parasitic or bacterial load that is systemic or localized in the digestive system.

SHOP NOW


More LSF-2 Product Details

Some pathogens associated with Lyme Disease are bacteria or parasitic in nature but not in the Lyme pathogen category. Immune system and detoxification support is essential in some cases where Lyme Disease is mixed with other pathogens unrelated to Lyme Disease. This formula can be used as a stand-alone or in combination with other LSF products.

Support for mitochondria and ammonia toxins are also recommended in anyone suffering with acute or chronic Lyme Disease









*These statements have not been evaluated by the Food and Drug Administration. This product(s) is not intended to diagnose, treat, cure, or prevent any disease.

*Check with your healthcare practitioner if you are nursing, pregnant, under 18 years of age, or following a doctor-prescribed protocol



Warning:

Consult a healthcare practitioner if you are pregnant or nursing, have a serious medical condition or use any medications. Keep out of the reach of children. Do not use if the safety seal is damaged or missing, store in a cool dry place.

LSF-2 Supplemental Facts

Serving size 2.5 ml Servings Per Container: 4 fl. oz (48) / 8 fl. oz.(96) Amount per serving % Daily Value* Proprietary Blend............ 2.5 mL *

LSF-2 Ingredients:

Artemisia Annua, Andrographis, Boneset, Japanese Knotweed, Milk Thistle Seed, Pau d'Arco Bark, Red Root, Shepherds Purse, Teasel Root; Worm-wood, Yellow Dock Root.

Other Ingredients: Certified Organic Cane Alcohol (40%), Purified Water

LSF-2 Suggested Use

Dosage: Dilute in water, 2.5ml (1/2) tsp for each 50 lb of weight up to 250lb or 2.5 tsp in one serving, 2 times daily until gone or as directed by your health care practitioner. Take on an empty stomach or 15 minutes before meals.

Dosage Chart

Dosage Chart: 50lb - 1/2 tsp 100lb -1 tsp 150lb - 1 1/2 tsp

200lb - 2 tsp

